

ANNEX 2 - ICD 10 CODES for TB Cases

Anatomical Site	Diagnostic Criteria	Definition of Terms		ICD 10 Code/s
Pulmonary (PTB)	Bacteriologically confirmed	Smear-positive	A patient with at least one (1) sputum specimen positive for AFB, with or without radiographic abnormalities consistent with active TB	A 15.0
		Culture-positive	A patient with positive sputum culture for MTB complex, with or without radiographic abnormalities consistent with active TB	A 15.1 <i>Note: if confirmed by both culture and smear, the code should be A15.0</i>
		Rapid Diagnostic test-positive	A patient with sputum positive for MTB complex using rapid diagnostic modalities such as Xpert MTB/RIF, with or without radiographic abnormalities consistent with active TB	A 15.0
	Clinically diagnosed	<p>A patient with two (2) sputum specimens negative for AFB or MTB or with smear not done due to specified conditions but with radiographic abnormalities consistent with active TB; <b>and</b> there has been no response to a course of empiric antibiotics and/or symptomatic medications; <b>and</b> who has been decided (either by the TBDC and/or physician) to have TB disease activity requiring a full course of anti-TB chemotherapy</p> <p><b>OR</b></p> <p>A child (less than 15 years old) with two (2) sputum specimens negative for AFB or with smear not done, who fulfills three (3) of the five (5) criteria for disease activity namely: signs and symptoms suggestive of TB; exposure to an active TB case; positive tuberculin test; abnormal chest radiograph suggestive of TB; and other laboratory findings suggestive of tuberculosis <b>and</b> who has been decided (either by the physician and/or TBDC) to have TB disease requiring a full course of anti-TB chemotherapy</p> <p><b>OR</b></p> <p>A patient with laboratory or strong clinical evidence for HIV/AIDS with two (2)</p>		<p>A16.0 (<i>if smear or MTB culture was done but negative</i>)</p> <p>A16.1 (<i>if smear or culture was not done</i>)</p> <p>For cases of clinically diagnosed PTB resulting from HIV, the following codes shall be used:</p> <p>B20.0, A16.0 (<i>if smear or culture negative</i>)</p> <p>B20.0, A16.1 (<i>if smear or culture</i></p>

Anatomical Site	Diagnostic Criteria	Definition of Terms	ICD 10 Code/s
		sputum specimens negative for AFB or MTB or with smear not done due to specified conditions but who, regardless of radiographic results, has been decided (either by physician and/or TBDC) to have TB disease activity requiring a full course of anti-TB chemotherapy.	<i>was not done)</i> B20.0, A16.2 ( <i>without mention of smear or culture confirmation</i> )
Extra-pulmonary (EPTB)	Bacteriologically confirmed	A patient with a smear/culture/new diagnostic test from a biological specimen in an extra-pulmonary site (i.e., organs other than the lungs) positive for AFB or MTB complex	A15.4 – A15.6, A15.8 <i>Note: 4<sup>th</sup> character of ICD 10 Code depends on the site</i>
	Clinically diagnosed	A patient with histological and/or clinical or radiologic evidence consistent with active extra-pulmonary TB and there is a decision by a physician to treat the patient with anti-TB drugs	A16.3 – A16.6, A16.8, A17 – A19