Annex F: Clinical and Social Criteria based on Joint Administrative Order

ANNEX A. COVID-19 Patient Algorithm for Triage and Hospitalization (C-PATH).

Note: The DOH may henceforth release an updated version, which shall be used for this Order.

FIGURE 1A. CLASSIFICATION OF CASES

Version 06 April 2020 (original)





FIGURE 1B. CONTACT TRACING PROTOCOL

Version 06 April 2020 (original)



FOOTNOTES

Exposure by travel

Travel from a country/area where there is sustained community level transmission

Exposure by residence

Lives in an LGU where there is sustained community level transmission

Exposure by contact

- Providing direct care to suspect, probable, or confirmed COVID-19 patients without using proper PPE (i.e. healthcare workers);
- Face-to-face contact with a probable or confirmed case within 1 meter and for more than 15 minutes;
- 3. Direct physical contact with a probable or confirmed case; OR
- 4. Other situations as indicated by local risk assessments

dAcute Respiratory Illness (ARI)

Flu-like symptoms (cough, colds, sore throat, body malaise; fatigue, fever)

Home Quarantine – All members of the household (including pets) must strictly stay at home


BHERT Monitoring

Barangay Health Emergency Response Team (BHERT)

- Accomplish a Case Identification Form (CIF)
- Ensure monitoring throughout the duration of isolation & guarantine
- Facilitate home care and basic needs
- A daily report shall be forwarded to the Municipality/City Epidemiology and Surveillance Units (MESU/CESU) which in turn are forwarded to the Provincial Epidemiology and Surveillance Units (PESU)

FIGURE 1C. COMMUNITY QUARANTINE PROTOCOL

Version 05 April 2020 (original)



FOOTNOTES

- Self-isolation strict isolation of the patient in a separate room or area in the household
- Home Quarantine All members of the household (including pets) must strictly stay at home

Situations where home guarantine is difficult

- 1. Living with vulnerable person (with comorbids or >60y/o)
- 2. No separate bedroom or bed not >1m away
- 3. No separate bathroom for patient
- 4. Not well-ventilated
- 5. No separate utensils and personal things
- 6. No separate towels for handwashing

BHERT Monitoring

Barangay Health Emergency Response Team (BHERT)

- Accomplish a Case Identification Form (CIF)
- Ensure monitoring throughout the duration of isolation & quarantine
- Facilitate home care and basic needs
- A daily report shall be forwarded to the Municipality/City Epidemiology and Surveillance Units (MESU/CESU) which in turn are forwarded to the Provincial Epidemiology and Surveillance Units (PESU)

Repeat Test Negative

 Two consecutive negative tests 24 hours apart is preferred or at least one negative test prior to discharge

Improvement of symptoms:

- Temp <37.8°C > 3 days,
- Respiratory symptoms reduced significantly
- CXR shows significant improvement