

**Annex A**  
**Disease severity classification of patients managed as**  
**COVID-19 and corresponding Intermediate Package**

<b>Classification</b>	<b>Manifestations</b>	<b>Intermediate Package (RT-PCR Test Negative)</b>
<b>Mild</b>	Fever, cough, fatigue, anorexia, myalgias  Other non-specific symptoms such as sore throat, nasal congestion, headache, diarrhea, nausea and vomiting, Loss of smell (anosmia) or loss of taste (ageusia) preceding the onset of respiratory symptoms  NO signs of pneumonia or hypoxia	None
<b>Moderate</b>	Fever, cough, dyspnea  With <b>non-severe pneumonia</b> : RR 21-30/minute, SpO <sub>2</sub> >92% on room air  <b>60 years old or more, OR with co-morbidity</b>	Intermediate Package for moderate pneumonia  Package Code: IMP01
<b>Severe</b>	Fever, cough, dyspnea  With severe pneumonia or severe acute respiratory infection: RR >30 breaths/minute, SpO <sub>2</sub> <92% on room air	Intermediate Package for severe/critical pneumonia  Package Code: IMP02
<b>Critical</b>	Onset within 1 week of known clinical insult (pneumonia) or new or worsening respiratory symptoms, progressing infiltrates on chest x-ray (CXR) or chest CT, with respiratory failure not fully explained by cardiac failure or fluid overload	
	Sepsis: life-threatening organ dysfunction caused by a dysregulated host response to suspected or proven infection, Signs of organ dysfunction: altered mental status, difficult or fast breathing, low oxygen saturation, reduced urine output, fast heart rate, weak pulse, cold extremities or low blood pressure, skin mottling, or laboratory evidence of coagulopathy, thrombocytopenia, acidosis, high lactate, or hyperbilirubinemia	
	Septic Shock: persisting hypotension despite volume resuscitation, requiring vasopressors to maintain MAP $\geq$ 65 mmHg and serum lactate level > 2 mmol/L	

References: Interim Guidance on the Clinical Management of Adult Patients with Suspected or Confirmed COVID-19 Infection Versions 2.1 and 3.1