

Annex A - Disease severity classification of patients managed as COVID-19 and corresponding Intermediate Package

Classification	Manifestations	Intermediate Package (RT-PCR Test Negative)
Mild	Fever, cough, fatigue, anorexia, myalgias Other non-specific symptoms such as sore throat, nasal congestion, headache, diarrhea, nausea and vomiting, Loss of smell (anosmia) or loss of taste (ageusia) preceding the onset of respiratory symptoms NO signs of pneumonia or hypoxia	None
Moderate	Fever, cough, dyspnea With non-severe pneumonia : RR 21-30/minute, SpO2 >92% on room air 60 years old or more, OR with co-morbidity	Intermediate Package for moderate pneumonia Package Code: IMP01
Severe	Fever, cough, dyspnea With severe pneumonia or severe acute respiratory infection: RR >30 breaths/minute, SpO2 <92% on room air	Intermediate Package for severe/critical pneumonia Package Code: IMP02
Critical	Onset within 1 week of known clinical insult (pneumonia) or new or worsening respiratory symptoms, progressing infiltrates on chest x-ray (CXR) or chest CT, with respiratory failure not fully explained by cardiac failure or fluid overload or <i>acute respiratory distress syndrome</i> Sepsis: life-threatening organ dysfunction caused by a dysregulated host response to suspected or proven infection, Signs of organ dysfunction: altered mental status, difficult or fast breathing low oxygen saturation, reduced urine output, fast heart rate, weak pulse, cold extremities or low blood pressure, skin mottling, or laboratory evidence of coagulopathy, thrombocytopenia, acidosis, high lactate, or hyperbilirubinemia Septic Shock: persisting hypotension despite volume resuscitation, requiring vasopressors to maintain MAP \geq 65 mmHg and serum lactate level >2 mmol/L	

References: Interim Guidance on the Clinical Management of Adult Patients with Suspected or Clinical Management of Adult Patients with Suspected or Confirmed COVID-19 Infection Versions 2.1 and 3.1

